DATE: Monday, October 07, 2002

Set Name Query			Hit Count Set Name result set	
DR = II	SPT_PGPB.JPAB.DWPI; PLUR=YES; OP=ADJ			
)D U	1,1 01 1,1 01 1, 1 1, 1 1, 1 1, 1 1, 1	10	L3	
L3	(L1 and (transgen\$ or disrupt\$ or knockout)) AnD ((@pd > 20020308)!)	10	113	
		2	L2	
L2	(retina-specific nuclear receptor) AnD ((wpd > 20020308):)	_		
		39	L1	
Ll	(retina-specific nuclear receptor of KNK) Allo ((wpd > 20020300))	•		
	by side $DB = U$ L3	by side OB=USPT,PGPB,JPAB,DWPI; PLUR=YES; OP=ADJ L3 (L1 and (transgen\$ or disrupt\$ or knockout)) AnD ((@pd > 20020308)!)	by side OB=USPT,PGPB,JPAB,DWPI; PLUR=YES; OP=ADJ L3 (L1 and (transgen\$ or disrupt\$ or knockout)) AnD ((@pd > 20020308)!) L2 (retina-specific nuclear receptor) AnD ((@pd > 20020308)!) 2	

END OF SEARCH HISTORY

```
$%^STN;HighlightOn= ***;HighlightOff=*** ;
 Trying 3106016892...Open
 Welcome to STN International! Enter x:x
 LOGINID:ssspta1633cxq
 TERMINAL (ENTER 1, 2, 3, OR ?):2
 ******** Welcome to STN International
                                  Web Page URLs for STN Seminar Schedule - N. America
   NEWS 1
  NEWS 2 Sep 17 IMSworld Pharmaceutical Company Directory name change to PHARMASEARCH
   NEWS 3 Oct 09 Korean abstracts now included in Derwent World Patents
                          Index
   NEWS 4 Oct 09 Number of Derwent World Patents Index updates increased NEWS 5 Oct 15 Calculated properties now in the REGISTRY/ZREGISTRY File NEWS 6 Oct 22 Over 1 million reactions added to CASREACT
   NEWS 7 Oct 22 DGENE GETSIM has been improved NEWS 8 Oct 29 AAASD no longer available
    NEWS 9 Nov 19 New Search Capabilities USPATFULL and USPAT2
NEWS 10 Nov 19 TOXCENTER(SM) - new toxicology file now available on STN
   NEWS 10 Nov 29 COPPERLIT now available on STN
NEWS 12 Nov 29 DWPI revisions to NTIS and US Provisional Numbers
NEWS 13 Nov 30 Files VETU and VETB to have open access
NEWS 14 Dec 10 WPINDEXWPIDSWPIX New and Revised Manual Codes for
   2002
NEWS 15 Dec 10 DGENE BLAST Homology Search
NEWS 16 Dec 17 WELDASEARCH now available on STN
NEWS 17 Dec 17 STANDARDS now available on STN
NEWS 18 Dec 17 New fields for DPCI
NEWS 19 Dec 19 CAS Roles modified
NEWS 20 Dec 19 1907-1946 data and page images added to CA and CAplus
NEWS 21 Jan 25 BLAST(R) searching in REGISTRY available in STN on the
   2002
     NEWS 22 Jan 25 Searching with the P indicator for Preparations
NEWS 23 Jan 29 FSTA has been reloaded and moves to weekly updates
NEWS 24 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update
                           frequency
      NEWS 25 Feb 19 Access via Tymnet and SprintNet Eliminated Effective 3/31/02
      NEWS 26 Mar 08 Gene Names now available in BIOSIS
      NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),
AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002
      NEWS HOURS STN Operating Hours Plus Help Desk Availability
      NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
       NEWS PHONE
NEWS PHONE
NEWS WWW
NEWS WW
NEWS W
NEWS WW
NEWS W
NEWS WW
NEWS W
NEWS
      Enter NEWS followed by the item number or name to see news on that
      specific topic.
         All use of STN is subject to the provisions of the STN Customer
         agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation
         of commercial gateways or other similar uses is prohibited and may
         result in loss of user privileges and other penalties.
         AB
       FILE 'HOME' ENTERED AT 15:15:05 ON 08 MAR 2002
       => FIL BIOSIS EMBASE CAPLUS
                                                                                                    SINCE FILE TOTAL
       COST IN U.S. DOLLARS
                                                                               ENTRY SESSION
       FULL ESTIMATED COST
       FILE 'BIOSIS' ENTERED AT 15:15:19 ON 08 MAR 2002
       COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)
        FILE 'EMBASE' ENTERED AT 15:15:19 ON 08 MAR 2002
        COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.
       FILE 'CAPLUS' ENTERED AT 15:15:19 ON 08 MAR 2002
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)
         => s retina-specific nuclear protein
L1 3 RETINA-SPECIFIC NUCLEAR PROTEIN
         PROCESSING COMPLETED FOR L1
                              1 DUP REM L1 (2 DUPLICATES REMOVED)
         => d bib abs 1-
YOU HAVE REQUESTED DATA FROM 1 ANSWERS - CONTINUE? Y/(N):y
```

```
L2 ANSWER 1 OF 1 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE 1
 AN 1991:318007 BIOSIS
UN BA92:28522
TI CHARACTERIZATION OF DEVELOPMENTALLY REGULATED AND
***RETINA***

****SPECIFIC***

****NUCLEAR***

****PROTEIN***

BINDING TO A SITE IN
THE UPSTREAM REGION OF THE RAT OPSIN GENE.
 DN BA92:28522
 AU MORABITO M A; YU X; BARNSTABLE C J
CS DEP. OPHTHALMOL. VISUAL SCI., YALE UNIV. SCH. MED., 330 CEDAR
 ST. NEW
 HAVEN, CONN. 06510.
SO J BIOL CHEM, (1991) 266 (15), 9667-9672.
CODEN: JBCHA3. ISSN: 0021-9258.
  FS BA; OLD
 LA English

AB DNase I protection and gel retardation assays have identified a sequence
5' to the transcription start site of the rat opsin gene that interacts
with nuclear proteins from mammalian retinas but not from a variety of
other neural and non-neural tissues. Following sodium dodecyl
  LA English
        sulfate-polyacrylamide gel electrophoresis and transfer to nitrocellulose the protein(s) responsible for this binding were identified with an
       the protein(s) responsible for this binding were identified with an oligonucleotide probe and were found to migrate with an apparent molecular size of 40 kilodalitons. The binding complex eluted from fast protein liquid chromatography gel filtration as a peak centered at 100 kilodalitons, suggesting the presence of more than one subunit. Binding activity could be detected in postnatal day 1 retinal extracts and increased over the next 2 weeks of development, a time course coincident with opsin gene expression and maturation of rod photoreceptors. Synthetic oligonucleotides with altered sequences showed that the binding was dependent upon residues in a CTAAT motif and was facilitated by surrounding GGCCCC sequences. The specificity of the binding interaction was measured by inhibition of complex formation in a gel retardation
         was measured by inhibition of complex formation in a gel retardation assay. The unaltered sequence was over 2 orders of magnitude more
          effective at inhibiting complex formation than either an unrelated DNA
         sequence or a concensus sequence corresponding to a known CCAAT box binding protein NF1.
    => s retina-specific nuclear receptor
L3 4 RETINA-SPECIFIC NUCLEAR RECEPTOR
    => dup rem 13
PROCESSING COMPLETED FOR L3
                       2 DUP REM L3 (2 DUPLICATES REMOVED)
    YOU HAVE REQUESTED DATA FROM 2 ANSWERS - CONTINUE? Y/(N):y
     L4 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2002 ACS
AN 2000:659546 CAPLUS
             134:362020
           Assignment of the NR2E3 gene to mouse chromosome 9 and to human
     chromosome
     15q22.33.fwdarw.q23
AU Rendtorff, N. D.; Vissing, H.; Tumer, Z.; Silahtaroglu, A.; Tommerup, N. CS Department of Medical Genetics, The Panum Institute, Copenhagen, Den. Cytogenet. Cell Genet. (2000), 89(3-4), 279-280
           CODEN: CGCGBR; ISSN: 0301-0171
     PB S. Karger AG
DT Journal
              The human and mouse NR2E3 gene (also known as PNR) encoding a ***retina*** - ***specific*** ***nuclear*** ***receptor*** was
            recently identified and found to be a ligand-dependent transcription factor. Here we report the mapping of the mouse Nr2e3 gene using radiation hybrid mapping, and refine the localization of the human NR2E3 gene using fluorescence in situ hybridization and radiation hybrid
             mapping. The mouse Nr2e3 gene was mapped to chromosome 9 between
       markers
            D9Mit102 and D9Mit207, while the human NR2E3 gene mapped to
       chromosome
       15q22.33.fwdarw.q23.
RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS
        RECORD
                        ALL CITATIONS AVAILABLE IN THE RE FORMAT
        L4 ANSWER 2 OF 2 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS
        INCIDUPLICATE 1
        AN 2000:110993 BIOSIS
        DN PREV200000110993
TI ***Retina*** - ***specific***
                                                                                                       ***receptor*
                                                                       ***nuclear***
              A potential regulator of cellular retinaldehyde-binding protein expressed
        A potential regulator of ceilular retinalonyde-binding protein expressed in retinal pigment epithelium and Muller glial cells.

AU Chen, Fang (1); Figueroa, David J.; Marmorstein, Alan D.; Zhang, Qing; Petrukhin, Konstantin; Caskey, C. Thomas; Austin, Christopher P.

CS (1) Department of Bone Biology WP 26A-1000, Merck Research Laboratories,
         West Point, PA, 19488 USA
SO Proceedings of the National Academy of Sciences of the United States of America, (Dec. 21, 1999) Vol. 96, No. 26, pp. 15149-15154.
               ISSN: 0027-8424.
```

LA English

SL English

AB In an effort to identify nuclear receptors important in retinal disease, we screened a retina cDNA library for nuclear receptors. Here we describe the identification of a ***retina*** - ***specific***

nuclear ***receptor*** (RNR) from both human and mouse. Human RNR is a splice variant of the recently published photoreceptor cell-specific nuclear receptor (Kobayashi, M., Takezawa, S., Hara, K., Yu, R. T., Umesono, Y., Agata, K., Taniwaki, M., Yasuda, K. & Umesono, K. (1999) Proc. Natl. Acad. Sci. USA 96, 4814-4819) whereas the mouse RNR is a mouse ortholog. Northern blot and reverse transcription-PCR analyses of a mouse ornoiog. Normern out and reverse transcription-PCR analyses of human mRNA samples demonstrate that RNR is expressed exclusively in the retina, with transcripts of apprxeq7.5 kb, apprxeq3.0 kb, and apprxeq2.3 kb by Northern blot analysis. In situ hybridization with multiple probes on both primate and mouse eye sections demonstrates that RNR is expressed in the retinal pigment epithelium and in Muller glial cells. By using the Gal4 chimeric receptor/reporter cotransfection system, the ligand binding domain of RNR was found to repress transcriptional activity in the absence of exogenous ligand. Gel mobility shift assays revealed that RNR can interact with the promoter of the cellular retinaldehyde binding protein gene in the presence of retinoic acid receptor (RAR) and/or retinoid X receptor (RXR). These data raise the possibility that RNR acts to regulate the visual cycle through its interaction with cellular retinaldehyde binding protein and therefore may be a target for retinal diseases such as retinitis pigmentosa and age-related macular degeneration.

=> s retina-specific nuclear receptor or RNR L5 547 RETINA-SPECIFIC NUCLEAR RECEPTOR OR RNR

=> s I5 (10a) (knockout or transgen? or disrupt?)
L6 2 L5 (10A) (KNOCKOUT OR TRANSGEN? OR DISRUPT?)

=> dup rem 16 PROCESSING COMPLETED FOR L6 1 DUP REM L6 (1 DUPLICATE REMOVED)

=> d bib abs

L7 ANSWER 1 OF 1 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC. DUPLICATE 1

AN 2001:526761 BIOSIS

DN PREV200100526761 TI Mutational and structural analyses of the ribonucleotide reductase inhibitor Sml1 define its Rnr1 interaction domain whose inactivation

allows suppression of med 1 and rad53 lethality.

AU Zhao, Xiaolan; Georgieva, Bilyana; Chabes, Andrei; Domkin, Vladimir;
Ippel, Johannes H.; Schleucher, Jurgen; Wijmenga, Sybren; Thelander, Lars;

Rothstein, Rodney (1)
CS (1) Department of Genetics and Development, College of Physicians and Surgeons, Columbia University, 701 West 168th St., New York, NY, 10032:

rothstein@cuccfa.ccc.columbia.edu USA SO Molecular and Cellular Biology, (December, 2000) Vol. 20, No. 23, pp. 9076-9083. print. ISSN: 0270-7306.

DT Article

LA English

SL

St. English

AB In budding yeast, MEC1 and RAD53 are essential for cell growth. Previously we reported that mec1 or rad53 lethality is suppressed by removal of Sml1, a protein that binds to the large subunit of ribonucleotide reductase (Rnr1) and inhibits RNR activity. To understand further the relationship between this suppression and the Sml1-Rnr1 interaction, we randomly mutagenized the Sml1 open reading frame. Seven mutations were identified that did not affect protein expression levels but relieved mec1 and rad53 inviability. Interestingly, all seven mutations abolish the Sml1 interaction with Rnr1, suggesting that this interaction causes the lethality observed in mec1 and rad53 strains. The mutant residues all cluster within the 33 C-terminal amino acids of the 104-amino-acid-long English tethality observed in mechanic radios strains. The introduces air cluster within the 33 C-terminal amino acids of the 104-amino-acid-long Sml1 protein. Four of these residues reside within an alpha-helical structure that was revealed by nuclear magnetic resonance studies. Moreover, deletions encompassing the N-terminal half of Sml1 do not interfere with its "**RNR*** inhibitory activity. Finally, the seven sml1 mutations also ***disrupt** the interaction with yeast Rnr3 and human R1, suggesting a conserved binding mechanism between Sml1 and the large subunit of RNR from different species.

---Logging off of STN---

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS

SINCE FILE TOTAL **ENTRY**

FULL ESTIMATED COST

46.65 46.80 DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

CA SUBSCRIBER PRICE

ENTRY SESSION

-0.62

SINCE FILE

STN INTERNATIONAL LOGOFF AT 15:25:18 ON 08 MAR 2002